



<p><b>DUREZOL</b> <b>PREDNISOLONE</b></p> <p>Shake before use</p>	<p><b>One drop</b> <b>__2__ Times Daily</b></p>	<p><b>4 weeks from the date of surgery</b></p> <p><b>Last day:</b></p>
<p><b>PROLENSA</b> <b>ILEVRO</b> <b>KETOROLAC</b></p> <p>Shake before use</p>	<p><b>One Drop</b> <b>__1_ Times Daily</b></p>	<p><b>4 weeks from the date of surgery</b></p> <p><b>Last day:</b></p>
<p><b>BESIVANCE</b> <b>CIPROFLOXACIN</b></p> <p>Shake before use</p>	<p><b>One Drop</b> <b>__2__ Times Daily</b></p>	<p><b>1 week from the date of surgery</b></p> <p><b>Last day:</b></p>

## **Post-operative Instructions for patients who have had cataract surgery**

Use drops as prescribed

Wait 5 minutes in between instillation of multiple drops

Avoid rubbing the eye

Covering the eye is not necessary

Use supplied sunglasses if desired (not necessary)

Wait 5 days before swimming or using eye makeup

All usual activities including showering, cooking, watching T.V., reading, and exercise can be done after the first day

### **You can expect:**

**Tearing**

**Burning**

**Redness**

**Stinging from drops**

**Some floating spots in the vision**

**Mild discharge on the edge of the eyelids**

It is normal to have fluctuating vision and difficulty seeing certain things for example, reading, until you have fully healed and the possible need for glasses has been addressed (usually 3-4 weeks)

### **CALL THE OFFICE IMMEDIATELY IF YOU HAVE:**

**Extreme pain or worsening swelling**

**Significant discharge from the eye**

**Sudden or quick visual loss**

**718-460-1200**